Tromboguard® Haemostatic dressing for first aid and treatment of traumatic wounds

Stop bleeding in 3 minutes!

- Tromboguard® is haemostatic i.e. bleeding stemming dressing
- the haemostatic activity of the dressing is immediate and occurs within 3 minutes after application of the dressing to the wound – it was confirmed in clinical studies
- the dressing consists of 3 layers, each of which fulfils a different function:
  - the active layer of the dressing is a unique on the market combination of active ingredients (chitosan, alginates and silver), which are responsible for stopping the bleeding
  - the next layer is polyurethane foam, which has absorbent properties and keeps the wound moist
  - the outer layer is a polyurethane membrane that protects the wound against external factors
- the dressing may remain on the wound for a maximum of 24 hours
- more about the dressing in the information leaflet

Application:

- is used to stop traumatic bleeding (e.g. caused by traffic accidents or in the field)
- ideal for first aid kits, emergency medical services, and use on the battlefield


Types:

- sterile

Size and packing:

- sterile 10 cm x 10 cm, packed by 5 pcs
- other sizes available on request







MD

Sterile devices are certified by the Notified Body PCBC SA and bear the CE mark with number 1434

>>

©Copyright 2026 en.matopat-global.com | Last updated: 04.11.2026